


BAHADUR MEMORIAL LECTURES

In honor of Raj Bahadur's fundamental contributions to statistics and to our department.



The University of Chicago, Department of Statistics, presents the
Nineteenth Annual Bahadur Memorial Lectures



Håvard Rue

King Abdullah University of Science and Technology

Monday, April 16, 2018

4:30 PM, Eckhart 133, 5734 S. University Avenue

Reception before lecture in Jones 111

“Some New Developments in the R-INLA Project”

I will discuss some recent developments and works in progress related to the R-INLA project (www.r-inla.org), which offers approximate Bayesian inference for latent Gaussian models and is extensively used by researchers worldwide. The topics will include quantile regression for discrete responses, an approximate representation of the fractional Gaussian noise model with linear computational complexity, Student-t response and random effects, and how to construct Gaussian fields which obeys physical barriers like islands and coastlines.

Thursday, April 19, 2018

3:30 PM, Stevanovich Center, MS 112, 5727 S. University Avenue

Reception following lecture in Stevanovich Center, MS Library

“Penalizing Model Component Complexity: A Principled Practical Approach to Constructing Priors”

Setting prior distributions on model parameters is the act of characterizing the nature of our uncertainty and has proven a critical issue in applied Bayesian statistics. Although the prior distribution should ideally encode the users' uncertainty about the parameters, this level of knowledge transfer seems to be unattainable in practice and applied statisticians are forced to search for a “default” prior. Despite the development of objective priors, which are only available explicitly for a small number of highly restricted model classes, the applied statistician has few practical guidelines to follow when choosing the priors. An easy way out of this dilemma is to re-use prior choices of others, with an appropriate reference. I will introduce a new concept for constructing prior distributions. We exploit the natural nested structure inherent to many model components, which defines the model component to be a flexible extension of a base model. Proper priors are defined to penalize the complexity induced by deviating from the simpler base model and are formulated after the input of a user-defined *scaling* parameter for that model component, both in the univariate and the multivariate case. These priors are invariant to reparameterizations, have a natural connection to Jeffreys' priors, are designed to support Occam's razor and seem to have excellent robustness properties, all which are highly desirable and allow us to use this approach to define default prior distributions. I will demonstrate the appropriateness of this approach and how it can be applied in various situations, like random effect models, spline smoothing, disease mapping, Cox proportional hazard models with time-varying frailty, spatial Gaussian fields and multivariate probit models. Further, we show how to control the overall variance arising from many model components in hierarchical models.